



**SANDHILLS
CENTER**



Video head impulse testing

Clinical Policy ID: CCP.1359

Recent review date: 3/2022

Next review date: 7/2023

Policy contains: Dizziness; head impulse testing; vestibular function; vertigo.

This policy is a Sandhills Center Clinical Coverage Policy adopted from AmeriHealth Caritas of North Carolina. These clinical policies are used to assist with making coverage determinations. Sandhills Center's clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of "medically necessary," and the specific facts of the particular situation are considered by Sandhills Center when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. Sandhills Center clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients. Sandhills Center's clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, Sandhills Center will update its clinical policies as necessary. Sandhills Center clinical policies are not guarantees of payment.

Coverage policy

Video head impulse testing is clinically proven and, therefore, medically necessary to evaluate horizontal semicircular canal function, when all of the following criteria are met (Alhabib, 2017; American Speech-Language-Hearing Association, 2021; Bhattacharyya, 2017; Fife, 2017):

- The member has symptoms of a vestibular disorder (e.g., dizziness, vertigo, imbalance, falls without other medical explanation).
- History and clinical evaluation, including the standard head impulse test, is inconclusive.
- The test results will aid in determining the appropriate medical or surgical treatment for disorders of auditory, balance, and other neural systems.

Limitations

Video head impulse testing is not medically necessary as an initial, sole screening test for dizziness (Cohen, 2017; Mezzalana, 2017).

Alternative covered services

- Computerized dynamic posturography.
- Electronystagmography or videonystagmography.
- Caloric testing.
- Rotation testing.
- Qualitative vestibular assessment.
- Scleral search coil method.
- Specialty referral.
- Standard of care diagnostic testing (e.g., audiometry, visual acuity examination, radiography, and blood work).

Background

Dizziness (including vertigo) affects approximately 15% to 30% of the general population, and its prevalence rises with age. The most common causes of dizziness are vestibular disorders (Neuhauser, 2016; Piker, 2016). Congenital or acquired conditions resulting from disease or injury to the vestibular system may reduce vision and depth perception and alter balance and coordination (American Speech-Language-Hearing Association, 2021).

A battery of non-instrumented tests (direct observation) and instrumented tests using both static and dynamic measures may be performed at the bedside or in an office setting to discern the etiology of vestibular dysfunction (American Speech-Language-Hearing Association, 2021). Within the context of the individual's medical history, diagnostic testing is performed to differentiate peripheral and central etiologies based on response to visual fixation and the presence and direction of nystagmus, which is a vision condition characterized by involuntary, rapid, and repetitive horizontal, vertical, or circular movement of the eyes (American Academy of Ophthalmology, 2020).

Standard and video head impulse tests

The standard head impulse test, also known as a head thrust test, impulse rotational test, or vestibulo-ocular reflex test, is used to assess horizontal semicircular canal function (Halmagyi, 2017). During a standard, non-instrumented head impulse test, the clinician rotates the patient's head abruptly and unpredictably in the vertical axis of the head while the patient keeps a fixed gaze on a target. A patient with vestibular impairment will move his or her eyes with the head, forcing a corrective eye movement (saccade) at the end of each head impulse to return to the target. Results may demonstrate a problem with semicircular canal function on the same side to which the head was turned prior to the corrective refixation response.

The standard procedure is noninvasive, safe, easy, and uses stimuli in the physiological range of everyday head movement, but it lacks objective documentation. The video version images the eye using sophisticated eye tracking and head velocity transducers, providing an objective record of eye movement and head movement during head impulses.

The Food and Drug Administration gave the first approval to sell a video head impulse testing system in the United States to GN Otometrics, manufacturer of the ICS Impulse, in February 2013 (GN Otometrics North America, 2013).

Findings

We identified three systematic reviews (Abouzayd, 2017; Alhabib, 2017; Verbecque, 2017), four prospective non-randomized comparative studies (Cohen, 2017; Eza-Nunez, 2016; Mezzalira, 2017; Ross, 2016), one retrospective comparative study (Skoric, 2017), and three evidence-based guidelines (American Speech-Language-Hearing Association, 2021; Bhattacharyya, 2017; Fife, 2017) for this policy. Three Medicare Local Coverage Determinations also address medical necessity criteria for video head impulse testing (Centers for Medicare & Medicaid Services, 2019a; 2019b; 2021).

The comparative diagnostic efficacy of video head impulse testing and other vestibular function tests is based on retrospective and prospective nonrandomized cohort studies of low to moderate quality (Abouzayd, 2017; Alhabib, 2017; Verbecque, 2017). Caloric testing, cervical vestibular evoked myogenic potentials, un-instrumented head impulse tests, head shaking tests, and rotational chair tests were the most common comparators. Few studies compared video head impulse testing to the scleral search coil method, considered a gold standard for accurate measurement of high velocity eye movements but is cumbersome to perform and non-portable.

Results suggest video head impulse testing provides distinct and complementary diagnostic information in the work up of persons with vestibular complaints when the clinical examination is inconclusive (Abouzayd, 2017; Alhabib, 2017; Verbecque, 2017). Limited evidence suggests quantifying eye movement with video head impulse testing may improve the ability to detect unobservable, covert saccades that occur during head impulses compared to its non-quantitative counterpart (Alhabib, 2017; Cohen, 2017).

There is an absence of evidence-based guidelines regarding which diagnostic algorithms should be applied to which patients among clinically heterogeneous populations who present with dizziness. Evidence-based guidelines agree that quantitative vestibular tests may be useful in evaluating vestibular disorders that may not be evident from the history and clinical examination (American Speech-Language-Hearing Association, 2021; Bhattacharyya, 2017; Fife, 2017). The choice of tests should be based on a case-by-case strategy according to each patient's symptoms and their suspected origin, and equipment availability.

In 2019, we updated the references, and the policy ID was changed from CP# 09.01.16 to CCP.1161.

In 2020, no policy changes are warranted.

In 2021, we updated the references. No policy changes are warranted.

In 2022, we added several large reviews, including:

- A systematic review of seven studies (n = 235) compared older adults with and without cognitive impairment. Video head impulse testing was unable to correlate cognitive decline with vestibulo-ocular reflex measurements in different frequency ranges of the semicircular canals (Bosmans, 2021).
- A meta-analysis of five studies (n = 253) of persons with benign paroxysmal positional vertigo or healthy controls found that video head impulse testing was valuable in diagnosing the condition, by detecting video-ocular reflex gains of the lateral, posterior, and anterior semi-circular canals on the affected side (Elsherif, 2021).
- A systematic review of 16 studies (n = 933) of persons with vestibular neuritis and controls determined the ability of video head impulse testing to diagnose the condition is high. Authors conclude the test is a

useful complement or alternative to caloric and rotational tests as an indicator of lesions of vestibular canal functioning, especially at the time of onset (Manzari, 2021).

- A systematic review/meta-analysis of 11 studies (n = 2,670) of patients with chronic dizziness showed video head impulse testing altered results in 21% of patients, compared with 55% altered after (gold standard) caloric testing. While video head impulse does not substitute for caloric testing, authors declare it to be complementary in assessing patients with dizziness (Vallim, 2021).
- An inter-rater reliability review of 171 patients presenting with acute vestibular syndrome showed the video head impulse test was highly accurate in differentiating vestibular neuritis from stroke. Authors conclude the test may be a feasible and effective tool to screen patients for potentially underlying stroke in the emergency setting (Machner, 2021).

References

On December 29, 2021, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were “Head Impulse Test” (MeSH), “reflex/diagnosis” (MeSH), “Head impulse test,” “head thrust test,” “impulse rotational test,” and “vestibulo-ocular reflex test.” We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

Abouzayd M, Smith PF, Moreau S, Hitier M. What vestibular tests to choose in symptomatic patients after a cochlear implant? A systematic review and meta-analysis. *Eur Arch Otorhinolaryngol*. 2017;274(1):53-63. Doi: 10.1007/s00405-016-4007-4.

Alhabib SF, Saliba I. Video head impulse test: A review of the literature. *Eur Arch Otorhinolaryngol*. 2017;274(3):1215-1222. Doi: 10.1007/s00405-016-4157-4.

American Academy of Ophthalmology. What is nystagmus? <https://www.aao.org/eye-health/diseases/what-is-nystagmus>. Reviewed October 28, 2020.

American Speech-Language-Hearing Association. Balance system disorders (Practice Portal). www.asha.org/Practice-Portal/Clinical-Topics/Balance-System-Disorders/. Published 2021.

Bhattacharyya N, Gubbels SP, Schwartz SR, et al. Clinical practice guideline: Benign paroxysmal positional vertigo (Update). *Otolaryngol Head Neck Surg*. 2017;156(3_suppl):S1-S47. Doi: 10.1177/0194599816689667.

Bosmans J, Jorissen C, Gilles A, et al. Vestibular function in older adults with cognitive impairment: A systematic review. *Ear Hear*. 2021;42(5):1119-1126. Doi: 10.1097/AUD.0000000000001040.

Center for Medicare & Medicaid Services. L33966 Vestibular function tests. <https://www.cms.gov/medicare-coverage-database/view/lcd.aspx?lcdid=33966&ver=18&bc=0>. Revised October 1, 2019. (a)

Center for Medicare & Medicaid Services. L34537 Vestibular function testing. <https://www.cms.gov/medicare-coverage-database/new-search/search.aspx>. Revised September 9, 2021.

Center for Medicare & Medicaid Services. L35007 Vestibular and audiologic function studies. <https://www.cms.gov/medicare-coverage-database/new-search/search.aspx>. Revised October 17, 2019. (b)

Cohen HS, Stitz J, Sangi-Haghpeykar H, et al. Utility of quick oculomotor tests for screening the vestibular system in the subacute and chronic populations. *Acta Otolaryngol*. 2017:1-5. Doi: 10.1080/00016489.2017.1398838.

Elsherif M, Eldeeb D, Eldeeb M. Clinical significance of video head impulse test in benign paroxysmal positional vertigo: A meta-analysis. *Eur Arch Otorhinolaryngol*. 2021;278(12):4645-4651. Doi: 10.1007/s00405-021-06832-3.

Eza-Nuñez P, Farinas-Alvarez C, Fernandez NP. Comparison of three diagnostic tests in detecting vestibular deficit in patients with peripheral vestibulopathy. *J Laryngol Otol*. 2016;130(2):145-150. Doi: 10.1017/s0022215115003114.

Fife TD, Colebatch JG, Kerber KA, et al. Practice guideline: Cervical and ocular vestibular evoked myogenic potential testing: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2017;89(22):2288-2296. Doi: h10.1212/WNL.0000000000004690.

Halmagyi GM, Chen L, MacDougall HG, et al. The video head impulse test. *Front Neurol*. 2017;8:258. Doi: 10.3389/fneur.2017.00258.

Machner B, Erber K, Choi JH, Sprenger A, Helmchen C, Trillenber. A simple gain-based evaluation of the video head impulse test reliably detects normal vestibulo-ocular reflex indicative of stroke in patients with acute vestibular syndrome. *Front Neurol*. 2021;12:741859. Doi: 10.3389/fneur.2021.741859.

Manzari L, Princi AA, DeAngelis S, Tramontano M. Clinical value of the video head impulse test in patients with vestibular neuritis: A systematic review. *Eur Arch Otorhinolaryngol*. 2021;278(11):4155-4167. Doi: 10.1007/s00405-021-06803-8.

GN Otometrics North America. FDA Clears GN Otometrics' ICS Impulse for Video Head Impulse Testing. <https://hearingreview.com/hearing-products/fda-clears-gn-otometrics-ics-impulse-for-video-head-impulse-testing>. Published February 20, 2013.

Mezzalira R, Bittar RSM, do Carmo Bilecki-Stipsky MM, Brugnera C, Grasel SS. Sensitivity of caloric test and video head impulse as screening test for chronic vestibular complaints. *Clinics (Sao Paulo)*. 2017;72(8):469-473. Doi: 10.6061/clinics/2017(08)03.

Neuhauser HK. The epidemiology of dizziness and vertigo. *Handb Clin Neurol*. 2016;137:67-82. Doi: 10.1016/b978-0-444-63437-5.00005-4.

Piker EG, Schulz K, Parham K, et al. Variation in the use of vestibular diagnostic testing for patients presenting to otolaryngology clinics with dizziness. *Otolaryngol Head Neck Surg*. 2016;155(1):42-47. Doi: 10.1177/0194599816650173.

Ross LM, Helminski JO. Test-retest and interrater reliability of the video head impulse test in the pediatric population. *Otol Neurotol*. 2016; 37(5): 558-563. Doi: 10.1097/mao.0000000000001040.

Skoric MK, Adamec I, Pavicic T, et al. Vestibular evoked myogenic potentials and video head impulse test in patients with vertigo, dizziness and imbalance. *J Clin Neurosci*. 2017;39:216-220. Doi: 10.1016/j.jocn.2017.02.009.

Vallim MGB, Gabriel GP, Messalira R, Stoler G, Chone CT. Does the video head impulse test replace caloric testing in the assessment of patients with chronic dizziness? A systematic review and meta-analysis. *Braz J Otorhinolaryngol*. 2021;87(6):733-741. Doi: 10.1016/j.bjorl.2021.01.002.

Verbecque E, Marijnissen T, De Belder N, et al. Vestibular (dys)function in children with sensorineural hearing

loss: a systematic review. *Int J Audiol*. 2017;56(6):361-381. Doi: 10.1080/14992027.2017.1281444.

Policy updates

1/2018: initial review date and clinical policy effective date: 3/2018

1/2019: references updated. The policy ID was changed from CP# 09.01.16 to CCP.1161.

3/2020: references updated.

3/2021: Policy references updated.

3/2022: Policy references updated.