



**SANDHILLS
CENTER**



Home-use light box therapy for seasonal affective disorder

Clinical Policy ID: CCP.1460

Recent review date: 2/2022

Next review date: 6/2023

Policy contains: Light box; light therapy; seasonal affective disorder; winter depression.

This policy is a Sandhills Center Clinical Coverage Policy adopted from AmeriHealth Caritas of North Carolina. These clinical policies are used to assist with making coverage determinations. Sandhills Center's clinical policies are based on guidelines from established industry sources, such as the Centers for Medicare & Medicaid Services (CMS), state regulatory agencies, the American Medical Association (AMA), medical specialty professional societies, and peer-reviewed professional literature. These clinical policies along with other sources, such as plan benefits and state and federal laws and regulatory requirements, including any state- or plan-specific definition of "medically necessary," and the specific facts of the particular situation are considered by Sandhills Center when making coverage determinations. In the event of conflict between this clinical policy and plan benefits and/or state or federal laws and/or regulatory requirements, the plan benefits and/or state and federal laws and/or regulatory requirements shall control. Sandhills Center clinical policies are for informational purposes only and not intended as medical advice or to direct treatment. Physicians and other health care providers are solely responsible for the treatment decisions for their patients. Sandhills Center's clinical policies are reflective of evidence-based medicine at the time of review. As medical science evolves, Sandhills Center will update its clinical policies as necessary. Sandhills Center clinical policies are not guarantees of payment.

Coverage policy

Home-use light box therapy is clinically proven and, therefore, medically necessary for seasonal affective disorder (major depressive disorder with seasonal patterns) when the following criteria are met:

- The member is diagnosed with recurring depression (at least two years) during a period when light decreases, which also alleviates when light increases (National Institute of Mental Health, 2021).
- An evaluation and recommendation for light box therapy is made by a physician, for treatment of no longer than one hour, most often early in the morning (Kurlansik, 2012; National Health Service, 2018).

Limitations

No limitations were identified during the writing of this policy.

Alternative covered services

- Antidepressants, especially selective serotonin reuptake inhibitors.
- Psychotherapy.
- Cognitive behavioral therapy.

Background

Seasonal affective disorder is a type of depression characterized by its recurrent seasonal pattern, with symptoms lasting about 4 to 5 months per year (National Institute of Mental Health, 2021). Low levels of serotonin and elevated levels of melatonin in the brain may contribute to this condition, with symptoms including low energy, hypersomnia, overeating, weight gain, craving for carbohydrates, and social withdrawal (feel like “hibernating”).

While the precise causes of the disorder are not known, risk factors are well documented. These include being female (80% are women), family history of other types of depression, having depression or bipolar disorder, and being younger (versus older) adults. Geographic prevalence varies greatly; for example an estimated 1.4% of Florida residents have the disease, compared to 9.9% in Alaska (Rohan, 2013).

While reports of the disorder have been made for many years, Lewy (1980) was the first to note improvement in the condition after exposure to bright light. The term “seasonal affective disorder” was first reported in the literature in 1984 (Rosenthal, 1984).

A variety of light boxes are used for seasonal affective disorder. Each filters out the ultraviolet rays and requires 20 – 60 minutes of exposure to 10,000 lux of cool-white fluorescent light, an amount that is about 20 times greater than ordinary indoor lighting (National Institute of Mental Health, 2021). Caution is advised not to sit directly facing the box, and at a reasonable proximity, namely 16 to 24 inches away (Mayo Clinic, 2016).

Bright light works by stimulating cells in the retina that connect to the hypothalamus, a part of the brain that helps control circadian rhythms. Activating the hypothalamus at a certain time every day can restore a normal circadian rhythm that normal light in winter cannot for some individuals (Miller, 2012).

Findings

The American Academy of Family Physicians recommends light box therapy for seasonal affective disorder, noting that improvement is usually observed one to four weeks after treatment starts. The Academy also recommends cognitive behavioral therapy and antidepressants (Kurlansik, 2012).

The British National Health Service states that cognitive behavioral therapy, antidepressants, and light therapy are all acceptable treatments for seasonal affective disorder. Light therapy works best in the morning, but is not appropriate for persons sensitive to bright light, or for those taking antibiotics and antipsychotics that increase light sensitivity (National Health Service, 2018).

A Canadian panel, while acknowledging light therapy to be one of the major options for persons with seasonal affective disorder, could not recommend either light or drug therapy over the other due to a lack of evidence (British Columbia Drug and Poison Information Centre, 2017).

The American Psychiatric Association guideline states that use of light box therapy “might be used” to treat seasonal affective disorder, and “may hasten the response to treatment with antidepressant medication” (American Psychiatric Association, 2010).

A meta-analysis of 18 randomized controlled trials (n = 610 adults with seasonal affective disorder) compared bright light therapy to dim light or sham/low-density negative ion generators as controls. Bright light therapy was superior to controls for depression ratings and response to active treatment. Authors recommended higher-quality trials due to moderate heterogeneity and moderate/high risk of bias, plus small sample sizes in the current research (Prjek, 2020).

A systematic review/meta-analysis (n = 397) found improvements in depression scores for patients with seasonal affective disorder given light therapy plus placebo were greater than those given antidepressants plus placebo,

and light therapy yielded a significantly greater improvement at $P < .001$ than those given antidepressants plus placebo; and also was significantly more effective for non-seasonal depression ($P < .005$). Authors note that light therapy has been found effective, but is underused in clinical settings (Geoffroy, 2019).

A systematic review of 40 studies observed bright light therapy demonstrated improvements to phase-advance delayed rhythms, sleep-wake disorders, and mood symptoms, but could not conclude that treatment results in changes in circadian outcomes (Menculini, 2018).

A systematic review (43 articles) of light therapy found no adverse ocular effects in otherwise healthy, unmedicated persons using the treatment. Further study is warranted for those with pre-existing ocular abnormalities or elevated photosensitivity (Brouwer, 2017).

A meta-analysis of two studies on seasonal affective disorder showed that persons using light box therapy reached significant improvement levels at the second and third weeks after initiation, and at the endpoint of the trials (Martensson, 2015).

In a Cochrane review of six studies, two ($n = 146$) compared outcomes for the second-generation antidepressant fluoxetine with those of light therapy for seasonal affective disorder. Trends in depression were similar in both groups, in which about two-thirds of subjects improved (Thaler, 2011).

In 2022, we found no new relevant literature to add to the policy.

References

On December 20, 2021, we searched PubMed and the databases of the Cochrane Library, the U.K. National Health Services Centre for Reviews and Dissemination, the Agency for Healthcare Research and Quality, and the Centers for Medicare & Medicaid Services. Search terms were "light box," "light therapy," "seasonal affective disorder," and "winter depression." We included the best available evidence according to established evidence hierarchies (typically systematic reviews, meta-analyses, and full economic analyses, where available) and professional guidelines based on such evidence and clinical expertise.

American Psychiatric Association. Practice guideline for the treatment of patients with major depressive disorder, third edition.

https://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/mdd.pdf. Published October, 2010.

British Columbia Drug and Poison Information Centre. Seasonal affective disorder (SAD).

<http://www.dpic.org/article/professional/seasonal-affective-disorder-sad>. Published 2017.

Brouwer A, Nguyen H-T, Snoek FJ, et al. Light therapy: Is it safe for the eyes? *Acta Psychiatr Scand*. 2017;136(6):534-548. Doi: 10.1111/acps.12785.

Geoffroy PA, Schroder CM, Reynaud E, Bourgin P. Efficacy of light therapy versus antidepressant drugs, and of the combination versus monotherapy, in major depressive episodes: A systematic review and meta-analysis. *Sleep Med Rev*. 2019;48:101213. Doi: 10.1016/j.smr.2019.101213.

Kurlansik SL, Ibay AD. Seasonal affective disorder. *Am Fam Physician*. 2012;86(11):1037-1041.

<https://www.ncbi.nlm.nih.gov/pubmed/?term=Kurlansik+SL%2C+Ibay+AD>.

Lewy AJ, Wehr TA, Goodwin FK, Newsome DA, Markey SP. Light suppresses melatonin secretion in humans. *Science*. 1980;210(4475):1267-1269. Doi: 10.1126/science.7434030.

Martensson B, Pettersson A, Berglund L, Ekselius L. Bright white light therapy in depression: A critical review of the evidence. *J Affect Disord*. 2015;182:1-7. Doi: 10.1016/j.jad.2015.04.013.

Mayo Clinic. Seasonal affective disorder treatment: Choosing a light therapy box. <https://www.mayoclinic.org/diseases-conditions/seasonal-affective-disorder/in-depth/seasonal-affective-disorder-treatment/art-20048298>. Published March 16, 2016.

Menculini G, Verdolini N, Murru A, et al. Depressive mood and circadian rhythms disturbances as outcomes of seasonal affective disorder treatment: A systematic review. *J Affect Disord*. 2018;241:608-626. Doi: 10.1016/j.jad.2018.08.071.

Miller MC. Seasonal affective disorder: Bring on the light. Harvard Medical School. <https://www.health.harvard.edu/blog/seasonal-affective-disorder-bring-on-the-light-201212215663>. Published December 21, 2012.

National Health Service. Treatment: Seasonal affective disorder (SAD). <https://www.nhs.uk/conditions/seasonal-affective-disorder-sad/treatment/>. Last reviewed July 30, 2018.

National Institute of Mental Health. Seasonal affective disorder. <https://www.nimh.nih.gov/health/topics/seasonal-affective-disorder/index.shtml>. Undated.

Prjek E, Friedrich M-E, Cambioli L, et al. The efficacy of light therapy in the treatment of seasonal affective disorder: A meta-analysis of randomized controlled trials. *Psychother Psychosom*. 2020;89(1):17-24. Doi: 10.1159/000502891.

Rohan KJ, Evans M, Mahon JN, et al. Cognitive-behavioral therapy vs. light therapy for preventing winter depression recurrence: Study protocol for a randomized controlled trial. *Trials*. 2013;14:82. Doi: 10.1186/1745-6215-14-82.

Rosenthal NE, Sack DA, Gillin JC, et al. Seasonal affective disorder. A description of the syndrome and preliminary findings with light therapy. *Arch Gen Psychiatry*. 1984;41(1):72-80. Doi: 10.1001/archpsyc.1984.01790120076010.

Thaler K, Delivuk M, Chapman A, Gaynes BN, Kaminski A, Gartlehner G. Second-generation antidepressants for seasonal affective disorder. *Cochrane Database Syst Rev*. 2011;(12):CD008591. Doi: 10.1002/14651858.CD008591.pub2.

Policy updates

5/2020: initial review date and clinical policy effective date: 6/2020

2/2021: Policy references updated.

2/2022: Policy references updated.